Table 2-2a

Runoff curve numbers for urban areas 1/

| Cover description | | Curve numbers for hydrologic soil group | | | | |
|---|-----------------------------|--|----------|----------|----------|--|
| _ | verage percent | | | - | | |
| | pervious area $\frac{2}{2}$ | А | В | С | D | |
| Fully developed urban areas (vegetation established) | | | | | | |
| Open space (lawns, parks, golf courses, cemeteries, etc.)⅔: | | | | | | |
| Poor condition (grass cover < 50%) | | 68 | 79 | 86 | 89 | |
| Fair condition (grass cover 50% to 75%) | | 49 | 69 | 79 | 84 | |
| Good condition (grass cover > 75%) | | 39 | 61 | 74 | 80 | |
| mpervious areas: | • | 00 | 01 | | 00 | |
| Paved parking lots, roofs, driveways, etc. | | | | | | |
| (excluding right-of-way) | | 98 | 98 | 98 | 98 | |
| Streets and roads: | • | 00 | 00 | 00 | 00 | |
| Paved; curbs and storm sewers (excluding | | | | | | |
| right-of-way) | | 98 | 98 | 98 | 98 | |
| Paved; open ditches (including right-of-way) | | 83 | 89 | 92 | 93 | |
| Gravel (including right-of-way) | | 76 | 85 | 89 | 91 | |
| Dirt (including right-of-way) | | 70 72 | 82 | 87 | 89 | |
| Vestern desert urban areas: | • | 12 | 02 | 01 | 00 | |
| Natural desert landscaping (pervious areas only) 4/ | | 63 | 77 | 85 | 88 | |
| Artificial desert landscaping (impervious areas only) = | • | 05 | | 00 | 00 | |
| desert shrub with 1- to 2-inch sand or gravel mulch | | | | | | |
| and basin borders) | | 96 | 96 | 96 | 96 | |
| Jrban districts: | • | 50 | 50 | 50 | 50 | |
| Commercial and business | . 85 | 89 | 92 | 94 | 95 | |
| Industrial | | 81 | 32 88 | 91 | 93 | |
| Residential districts by average lot size: | . 14 | 01 | 00 | 31 | 30 | |
| 1/8 acre or less (town houses) | . 65 | 77 | 85 | 90 | 92 | |
| 1/4 acre | | 61 | 75 | 83 | 92 87 | |
| 1/4 acre | | 57 | 73 72 | 81 | 86 | |
| 1/2 acre | | 54 | 70 | 80 | 85 | |
| 1/2 acre | | $54 \\ 51$ | 68 | 80 79 | 84 | |
| 2 acres | | 46 | 65 | 79 77 | 04 82 | |
| 2 acres | . 12 | 40 | 05 | " | 02 | |
| Developing urban areas | | | | | | |
| Newly graded areas | | | | | | |
| (pervious areas only, no vegetation) ^{5/} | | 77 | 86 | 91 | 94 | |
| dle lands (CN's are determined using cover types | | | | | | |
| | | | | | | |

similar to those in table 2-2c). ¹ Average runoff condition, and $I_a = 0.2S$.

² The average percent impervious area shown was used to develop the composite CN's. Other assumptions are as follows: impervious areas are directly connected to the drainage system, impervious areas have a CN of 98, and pervious areas are considered equivalent to open space in good hydrologic condition. CN's for other combinations of conditions may be computed using figure 2-3 or 2-4.

³ CN's shown are equivalent to those of pasture. Composite CN's may be computed for other combinations of open space

⁴ Composite CN's for natural desert landscaping should be computed using figures 2-3 or 2-4 based on the impervious area percentage (CN = 98) and the pervious area CN. The pervious area CN's are assumed equivalent to desert shrub in poor hydrologic condition.

⁵ Composite CN's to use for the design of temporary measures during grading and construction should be computed using figure 2-3 or 2-4 based on the degree of development (impervious area percentage) and the CN's for the newly graded pervious areas.

cover type.

Table 2-2bRunoff curve numbers for cultivated agricultural lands 1/2

| | Cover description | Curve numbers for | | | | | |
|--|----------------------------|-------------------------|-----------------------|----|----|----|--|
| | Cover description | Hydrologic | hydrologic soil group | | | | |
| Cover type | Treatment ^{2/} | condition $\frac{3}{2}$ | А | В | С | D | |
| Fallow | Bare soil | | 77 | 86 | 91 | 94 | |
| 1 allow | Crop residue cover (CR) | Poor | 76 | 85 | 90 | 93 | |
| | crop residue cover (OR) | Good | 74 | 83 | 88 | 90 | |
| Row crops | Straight row (SR) | Poor | 72 | 81 | 88 | 91 | |
| 1 | | Good | 67 | 78 | 85 | 89 | |
| | SR + CR | Poor | 71 | 80 | 87 | 90 | |
| | | Good | 64 | 75 | 82 | 85 | |
| | Contoured (C) | Poor | 70 | 79 | 84 | 88 | |
| | | Good | 65 | 75 | 82 | 86 | |
| | C + CR | Poor | 69 | 78 | 83 | 87 | |
| | | Good | 64 | 74 | 81 | 85 | |
| | Contoured & terraced (C&T) | Poor | 66 | 74 | 80 | 82 | |
| contourou a terracea (car) | Good | 62 | 71 | 78 | 81 | | |
| | C&T+ CR | Poor | 65 | 73 | 79 | 81 | |
| | Good | 61 | 70 | 77 | 80 | | |
| Small grain SR SR + CR C C + CR C&T C&T+ CR | SR | Poor | 65 | 76 | 84 | 88 | |
| | | Good | 63 | 75 | 83 | 87 | |
| | SR + CR | Poor | 64 | 75 | 83 | 86 | |
| | | Good | 60 | 72 | 80 | 84 | |
| | С | Poor | 63 | 74 | 82 | 85 | |
| | | Good | 61 | 73 | 81 | 84 | |
| | C + CR | Poor | 62 | 73 | 81 | 84 | |
| | | Good | 60 | 72 | 80 | 83 | |
| | Poor | 61 | 72 | 79 | 82 | | |
| | Good | 59 | 70 | 78 | 81 | | |
| | Poor | 60 | 71 | 78 | 81 | | |
| | Good | 58 | 69 | 77 | 80 | | |
| Close-seeded | SR | Poor | 66 | 77 | 85 | 89 | |
| or broadcast | | Good | 58 | 72 | 81 | 85 | |
| legumes or | С | Poor | 64 | 75 | 83 | 85 | |
| rotation | | Good | 55 | 69 | 78 | 83 | |
| meadow | C&T | Poor | 63 | 73 | 80 | 83 | |
| | | Good | 51 | 67 | 76 | 80 | |

 $^{\rm 1}$ Average runoff condition, and $\rm I_a{=}0.2S$

² Crop residue cover applies only if residue is on at least 5% of the surface throughout the year.

³ Hydraulic condition is based on combination factors that affect infiltration and runoff, including (a) density and canopy of vegetative areas, (b) amount of year-round cover, (c) amount of grass or close-seeded legumes, (d) percent of residue cover on the land surface (good \geq 20%), and (e) degree of surface roughness.

Poor: Factors impair infiltration and tend to increase runoff.

Good: Factors encourage average and better than average infiltration and tend to decrease runoff.

Table 2-2c

Runoff curve numbers for other agricultural lands $\underline{1}'$

| Cover description | | Curve numbers for hydrologic soil group | | | | |
|--|----------------------|--|----|-----|----|--|
| Cover type | Hydrologic condition | А | B | C C | D | |
| Pasture, grassland, or range—continuous | Poor | 68 | 79 | 86 | 89 | |
| forage for grazing. ^{2/} | Fair | 49 | 69 | 79 | 84 | |
| | Good | 39 | 61 | 74 | 80 | |
| Meadow—continuous grass, protected from grazing and generally mowed for hay. | — | 30 | 58 | 71 | 78 | |
| Brush—brush-weed-grass mixture with brush | Poor | 48 | 67 | 77 | 83 | |
| the major element. ${}^{\underline{y}}$ | Fair | 35 | 56 | 70 | 77 | |
| | Good | 30 4∕ | 48 | 65 | 73 | |
| Woods—grass combination (orchard | Poor | 57 | 73 | 82 | 86 | |
| or tree farm). 5/ | Fair | 43 | 65 | 76 | 82 | |
| | Good | 32 | 58 | 72 | 79 | |
| Woods. ^{6/} | Poor | 45 | 66 | 77 | 83 | |
| | Fair | 36 | 60 | 73 | 79 | |
| | Good | 30 4/ | 55 | 70 | 77 | |
| Farmsteads—buildings, lanes, driveways, and surrounding lots. | — | 59 | 74 | 82 | 86 | |

¹ Average runoff condition, and $I_a = 0.2S$.

² *Poor:* <50%) ground cover or heavily grazed with no mulch.

Fair: 50 to 75% ground cover and not heavily grazed.

Good: > 75% ground cover and lightly or only occasionally grazed.

Poor: <50% ground cover.

3

Fair: 50 to 75% ground cover.

Good: >75% ground cover.

 4 $\,$ Actual curve number is less than 30; use CN = 30 for runoff computations.

⁵ CN's shown were computed for areas with 50% woods and 50% grass (pasture) cover. Other combinations of conditions may be computed from the CN's for woods and pasture.

⁶ *Poor:* Forest litter, small trees, and brush are destroyed by heavy grazing or regular burning. *Fair:* Woods are grazed but not burned, and some forest litter covers the soil. *Good:* Woods are protected from grazing, and litter and brush adequately cover the soil.

Table 2-2d

Runoff curve numbers for arid and semiarid rangelands $1\!\!/$

| Cover description | | Curve numbers for hydrologic soil group | | | | |
|--|-------------------------|--|----|----|----|--|
| Cover type | Hydrologic condition 2/ | A 3⁄ | В | С | D | |
| Herbaceous-mixture of grass, weeds, and | Poor | | 80 | 87 | 93 | |
| low-growing brush, with brush the | Fair | | 71 | 81 | 89 | |
| minor element. | Good | | 62 | 74 | 85 | |
| Oak-aspen—mountain brush mixture of oak brush, | Poor | | 66 | 74 | 79 | |
| aspen, mountain mahogany, bitter brush, maple, | Fair | | 48 | 57 | 63 | |
| and other brush. | Good | | 30 | 41 | 48 | |
| Pinyon-juniper—pinyon, juniper, or both; | Poor | | 75 | 85 | 89 | |
| grass understory. | Fair | | 58 | 73 | 80 | |
| | Good | | 41 | 61 | 71 | |
| Sagebrush with grass understory. | Poor | | 67 | 80 | 85 | |
| | Fair | | 51 | 63 | 70 | |
| | Good | | 35 | 47 | 55 | |
| Desert shrub—major plants include saltbush, | Poor | 63 | 77 | 85 | 88 | |
| greasewood, creosotebush, blackbrush, bursage, | Fair | 55 | 72 | 81 | 86 | |
| palo verde, mesquite, and cactus. | Good | 49 | 68 | 79 | 84 | |

¹ Average runoff condition, and $I_{av} = 0.2S$. For range in humid regions, use table 2-2c.

 2 $\,$ Poor: $\,$ <30% ground cover (litter, grass, and brush overstory).

Fair: 30 to 70% ground cover.

Good: > 70% ground cover.

³ Curve numbers for group A have been developed only for desert shrub.